

# Shear instabilities in granular mixtures

Massimo Pica Ciamarra,<sup>\*</sup> Antonio Coniglio,<sup>†</sup> and Mario Nicodemi<sup>‡</sup>

*Dip.to di Scienze Fisiche, Università di Napoli "Federico II"*

*INFN-Coherentia, INFN and AMRA, Napoli, Italy*

(Dated: December 21, 2004)

Dynamical instabilities in fluid mechanics are responsible of a variety of important common phenomena, such as waves on the sea surface or Taylor vortices in Couette flow. In granular media dynamical instabilities has just begun to be discovered. Here we show by means of molecular dynamics simulation the existence of a new dynamical instability of a granular mixture under oscillating horizontal shear, which leads to the formation of a striped pattern where the components are segregated. We investigate the properties of such a Kelvin-Helmholtz like instability and show how it is connected to pattern formation in granular flow and segregation.

PACS numbers: 45.70.Qj, 45.70.Mg, 47.54.+r

A simple example of instabilities in fluid mechanics is the Kelvin-Helmholtz instability where a flat interface between two fluids flowing one past the other at different velocities is unstable, explaining for instance why waves on the sea surface form [1, 2]. In athermal systems such as granular mixtures, where an hydrodynamic like theory is not yet well established, the discovery of instabilities is a recent achievement [3, 4, 5, 6]. By studying the evolution of two bands of different grains placed on an horizontally oscillating tray (see Fig. 1(a)), we discover a new dynamical instability. This instability appears as growing wavy interface, which, at long times, leads to a pattern of alternating segregated stripes of grains, perpendicular to the driving direction. We suggest that this type of surface shear instability is the common mechanism for apparently different phenomena such wave formation as found in experiments on granular flow [6], and segregation processes [7] as observed in [8, 9, 10, 11], initially explained in terms of a thermodynamic driven phase separation. In our simulations we determine the region of existence of the pattern formation - segregation process as a function of the area fractions of large and small grains, and we discuss the dependence of the pattern as a function of the amplitude and frequency of oscillation.

*MD simulation model* – We perform Molecular Dynamics (MD) simulations (also known as Discrete Element Methods) of a binary system of disks which lay on a tray (i.e., in two dimensions) with periodic boundary conditions in the x-direction and hard walls in the other. Grains interact with the tray by a viscous force proportional to their relative velocity via a viscosity parameter  $\mu$  different for the two species; when overlapping, grains interact via 2D Hertzian contact forces [12]. Simulations details and parameters are chosen to model experimental conditions similar to those investigated by Mullin and coworkers [8, 9, 11, 13].

Two grains with diameters  $D_i$  and  $D_j$  in positions  $\mathbf{r}_i$  and  $\mathbf{r}_j$  interact if overlapping, i.e., if  $\delta_{ij} = [(D_i + D_j)/2 - |\mathbf{r}_i - \mathbf{r}_j|] > 0$ . The interaction is given by a normal Hertz force with viscous dissipation [14, 15]. In

2D this reduces to the linear spring-dashpot model,  $\mathbf{f}_n = k_n \delta_{ij} \mathbf{n}_{ij} - \gamma_n m_{\text{red}} \mathbf{v}_{nij}$ , where  $k_n$  and  $\gamma_n$  are the elastic and viscoelastic constants, and  $m_{\text{red}} = m_i m_j / (m_i + m_j)$  is the reduced mass. As in [13] we model the interaction with the tray via a viscous force  $\mathbf{f}_t = -\mu(\mathbf{v} - \mathbf{v}_{\text{tray}})$ , where  $\mathbf{v}_{\text{tray}}(t) = 2\pi A \nu \sin(\nu t) \mathbf{x}$  is the velocity of the tray and  $\mathbf{v}$  the velocity of the disk, plus a white noise force  $\xi(t)$  with  $\langle \xi(t) \xi(t') \rangle = 2\Gamma \delta(t - t')$ , where  $\Gamma = 0.2 \text{ g}^2 \text{cm}^2 \text{s}^{-3}$ . The interaction between particles and walls is elastic. We solve the equations of motion by Verlet algorithm with an time-step of  $6\mu\text{s}$ . For the grain-grain interaction, we use the value  $k_n = 2 \cdot 10^5 \text{ g cm}^2 \text{s}^{-2}$  and  $\gamma_n$  chosen, for each kind of grains, such that the restitution coefficient is given:  $e = 0.8$  [14]. The two components of our mixture (named  $b$  and  $s$ ) have viscous coefficient  $\mu_b = 0.28 \text{ g s}^{-1}$  and  $\mu_s = 0.34 \text{ g s}^{-1}$ . Apart from a simple rescaling of masses and lengths, these values are those of Ref. [13] (and given in private communications), and are taken from direct measurements on the experimental system in [8]. The size of the tray is  $L_x = 320 \text{ cm}$ ,  $L_y = 16 \text{ cm}$ . The qualitative picture we discuss does not change if these values are changed.

*Results* – We consider by now a mixture of heavy grains, of mass  $M_b = 1 \text{ g}$  and area fraction  $\phi_b = 0.37$ , and light grains, with  $M_s = 0.03 \text{ g}$  and  $\phi_s = 0.41$ , all with the same diameter  $D = 1 \text{ cm}$ , prepared in a horizontally fully segregated configuration (Fig. 1(a)). The mixture is vibrated on an horizontal tray along the x-direction with amplitude  $A$  and frequency  $\nu$ . The characteristic time scale,  $\tau_b = M_b/\mu_b$  and  $\tau_s = M_s/\mu_s$ , is different for the two species and they are forced to oscillate with different amplitudes and phases, subject to an effective surface shear periodically varying in time. One may expect configurations in which the two different components form two stripes parallel to the driving direction (see Fig. 1(a)) to be stable, the two stripes oscillating independently with different amplitude and phases. Instead the initially flat interface between the two components is observed to evolve via the formation of a surface modulation which has a growing amplitude.




FIG. 1: (color on-line) Evolution of a binary mixture of heavy (red-dark gray) and light (blue-ligh gray) grains on an horizontal tray oscillating along the  $x$ -direction with amplitude  $A = 1.2$  cm and frequency  $\nu = 12$  Hz (only a quarter of the system length is shown in each figure). **Panels a** The system is initially prepared in a horizontally segregated configuration with a flat interface between the two species. Under shaking it develops a surface instability with a sine like modulation of growing amplitude leading at long times to a state where segregated stripes perpendicular to the driving direction appear. **Upper small panels** show the corresponding evolution of the surface instability model Eq.s (1) in presence of an oscillating shear  $\Delta v(t) = 2\pi A\nu \sin(2\pi\nu t)$ . **Panels b** The same system is now initially prepared in a mixed state and then shaken. The formation of local fluctuations of the density of the two species (shown at  $t = 1.6$  s) leads to microspheres generating the same instability mechanism as in panels a.

Finally, this leads to a state where segregated stripes appear, perpendicular to the driving direction (at variance with known results in colloidal fluids [16]). This instability appears of the same kind of that observed in two fluid systems and in liquid sand systems subject to horizontal oscillations under gravity, which is responsible for the ripples observed on the shoreline [17]. In these cases, however, because of gravity there is an energy cost associated to the growth of the interface, which therefore gets stabilized.

When the system starts from a disordered initial state, as shown in Fig. 1(b), the instability develops too. The formation of local fluctuations of the density of the two species create microspheres which are forced to evolve according to the instability shown in Fig. 1(a). For instance Fig. 1(b), at  $t = 1.6$  s, shows the existence of clusters of particles of the same species elongated perpendicularly to the driving direction. As time goes on these clusters merge giving rise to the striped pattern of Fig. 1(b) ( $t = 80$  s). It is apparent that the originally disordered system is ordered, i.e., segregated, by the dynamical instability. Fig. 1(a) ( $t = 320$  s) and Fig. 1(b) ( $t = 80$  s) show that the wavelength of the final steady state depends on the initial conditions.

The segregation process shown in Fig. 1(a) reproduces that observed experimentally in [8, 11]. In the present set-up grains radii are equal, no “depletion” forces can be at work, and no “thermodynamic” coarsening phenomena can be considered responsible for this segregation process, as initially conjectured [11] by making an analogy with colloidal systems. Such a remark appears to be consistent with a scenario proposed by use of modified Navier-Stokes equations [18]. Interestingly, segregation from a disordered initial state to a final striped one

has also been observed in a two fluid system oscillated in absence of gravity [19].

The above dynamical instabilities should manifest in the mixture also under different shearing protocols. To check this, we study now a different configuration where gravity induces granular flow and thus a differential shear on the mixture components appears. The tray no longer oscillates, but it is inclined of an angle  $\theta$  with the respect to the horizontal by rotation around the  $y$ -axis. The two species of grains attain different limit velocities (of the order of  $v_b \simeq g\tau_b \sin \theta$  and  $v_s \simeq g\tau_s \sin \theta$ , with  $g$  gravitational acceleration) and thus experience a differential shear. As shown in Fig. 2, under these conditions the initially flat interface between the mixture components changes too, via Kelvin-Helmholtz-like instabilities. An unstable pattern forms and curl-like structures appear. This phenomenon, induced by surface dynamical instabilities, is very similar to those recently discovered in experiments on granular flows [6].

The results of Fig. 1 and Fig. 2 point out the essential role of shear induced surface instabilities in pattern formation and the connections with segregation transitions. Even though this is not an usual fluid, the complex nature of this phenomenon can be understood via schematic hydrodynamic-like considerations. On the two sides of the perturbed interface flux lines are narrowed and widened as the pressure is respectively lowered and increased (Bernoulli law). The pressure gradient reinforces the perturbation of the interface giving rise to a positive feedback. Within such a perspective, the essential features of pattern formation observed in the two experiments of Figs 1 and 2 are captured by a model originally proposed for the Kelvin-Helmholtz instability of granular flows [6]. The model describes the evolution




FIG. 2: (color on-line) Evolving interface in the same granular mixture of Fig. 1a flowing down a tray rotated of an angle  $\theta = 12^\circ$  with respect to the horizontal around the y-axis. The red (dark gray) grains are faster than the blue (light gray) ones. **Upper small panels** The upper panels show the evolution, at corresponding times, of a Kelvin-Helmholtz-like surface dynamical instability modeled by Eq.s (1) in presence of a constant shear  $\Delta v(t) = \Delta v_0$ , reproducing a pattern qualitatively similar to the one found in the main panels.

of the velocity components,  $v_X$  and  $v_Y$ , of the system interface initially located at  $Y(X) = 0$ :

$$v_X = \Delta v(t) \tanh(Y) \quad v_Y = c \sin(k_X X) \quad (1)$$

where  $\Delta v(t)$  is the relative bulk velocity between grains far on the two side of the interface at time  $t$ , as  $k_X$  and  $c$  are constant parameters (we consider the same values used in [6], i.e.,  $k_X = 5$ ,  $c = 0.1$ , but our results are robust to changes). The presence of a velocity gradient,  $\Delta v(t)$ , independently from its microscopic origin (different grains friction, different driving of the components, etc...), enhances shear and results in patterns formation. In the case of gravity induced deformations,  $\Delta v(t) = \Delta v_0 \equiv v_b - v_s$  can be considered to be time independent since in the stationary regime the two limit velocities are given. In such a case, the model essentially coincides with the one introduced in [6] and well describes the qualitative features of the MD simulations of Fig. 2 (see upper panels). In the case of the vibrated horizontal tray, the relative bulk velocity is oscillating in time and we fix  $\Delta v(t) = 2\pi A\nu \sin(2\pi\nu t)$ . This gives rise (see upper panels in Fig. 1(a)) to a growing interface with features very close to those observed in the simulations.

*Pattern formation* – We show now under which conditions pattern formations occur and explain how the properties of the final state of the mixture, such as the emerging characteristic length scale of the stripes,  $\lambda$ , depend on both the dynamics control parameters and the relative concentration of the two components. This gives rise to a complex mixing/segregation diagram which we discuss in details here in the case of the horizontally vibrated experiment. In order to make direct comparisons with experimental results on segregation observed by Mullin and coworkers [8, 9, 11, 13], we consider now a binary mixture of large mono-disperse disks, of diameter  $D_b = 1\text{cm}$ , covering an area fraction  $\phi_b$ , and small poly-disperse disks [21], with average diameter  $D_s = 0.7\text{cm}$  (with 17% polydispersity) and area fraction  $\phi_s$ . Masses and frictional parameters are as given before.

The system starts from a mixed initial configuration as in Fig. 2(b). The nature of the state reached by the

system under shaking at late stage is crucially dependent on the area fraction of the two species: segregation in stripes is only found for high enough concentrations. This behaviour, in the  $(\phi_b, \phi_s)$  plane, is summarized in the diagram of Fig. 3(a) showing the system “fluid” and “crystal” regions along with their segregation properties, for  $\nu = 12\text{ Hz}$  and  $A = 1.2\text{ cm}$ . Large grains are considered to be in a “fluid” configuration when their radial density distribution function,  $g(r)$ , shows a first peak at  $r = D_b$  and a second one at  $r = 2D_b$ , and to be in a “crystal” configuration when a new peak at  $r = \sqrt{3}D_b$  appears [11]. The system is in a “glassy” state [22] when on the longest of our observation time scales, the system is still far from stationarity.

Fig. 3(a) shows that grains at small concentrations are mixed and in a fluid state. Segregation via stripes formation appears at higher concentrations. At even higher concentrations, large grains form stripes with a crystalline order, as smaller grains are always fluid for their polydispersity. Finally, at very high area fractions, the system is blocked in its starting disordered configuration (“glassy” region). For instance, by increasing  $\phi_s$  at a fixed value of  $\phi_b$  (say  $\phi_b \simeq 0.174$ ), we observe first a transition from a mixed fluid state to a segregated striped fluid and then a transition where the monodisperse phase crystallize. The experiments of [11], where  $\phi_b \simeq 0.174$ , show the very same transitions found here at locations differing by a 10%.

In the case  $\phi_b = 0.30$  and  $\phi_s = 0.28$ , where stripes form, we describe their dependence on the dynamics control parameters in Figs 3(b) and 3(c), showing that the length scale,  $\lambda$ , increases as a function of the shaking frequency,  $\nu$ , and of the amplitude  $A$ . These results are to be compared, for instance, with those found in liquid-sand mixtures under oscillating flow, where ripples form with a wavelength depending on the amplitude of oscillation, but not on its frequency [20]. The dependence on  $\nu$  can be schematically understood by comparison with the characteristic time scales  $\tau_b = M_b/\mu_b$  and  $\tau_s = M_s/\mu_s$  of the two species (here  $\tau_b^{-1} = 0.28\text{ Hz}$  and  $\tau_s^{-1} = 11.3\text{ Hz}$ ): in the limit  $\nu \gg \tau_b^{-1}, \tau_s^{-1}$  grains are not able to follow the tray motion and no sensitivity to  $\nu$  is expected, as




FIG. 3: **Panel a** Ordering properties of the late stage configurations of the mixture as a function of the area fractions of the two components. The *shaded area* covers the region where segregation via stripes formation occurs. **circles**: large grains are in a fluid state. **squares**: large grains form a crystal. **stars**: the system appears blocked in a “glassy” disordered configuration (see text). When stripes form their characteristic length scale  $\lambda$  is a function of the frequency,  $\nu$ , and of the amplitude,  $A$ , of the driving oscillations. This is shown, in the case  $\phi_b = 0.30$  and  $\phi_s = 0.28$ , in **panels b** and **c**.

well as when  $\nu \ll \tau_s^{-1}$  since the grains move with the tray. Analogously, the dependence on  $A$  is expected to be substantial when  $A$  is at least of the order of the mean grains separation length,  $l = (4\phi_b/\pi D_b^2 + 4\phi_s/\pi D_s^2)^{-1/2}$ , since under this condition grains strongly interact.

In conclusion our simulations of a binary mixture under horizontal vibrations have revealed the existence of a new dynamical instability and shed light on the process of size segregation under oscillatory shear and its connections to pattern formation in granular flows. Within such a unifying framework, we derived the “phase diagram” of the mixing/segregation states of the mixture and its corresponding transitions. Finally, such hydrodynamic like processes appear to be related to those known in thermal fluids and fluid-grains systems, ranging from Kelvin-Helmholtz instabilities to ripples formation in liquid sand mixtures, even though interesting difference are found.

Work supported by EU Network MRTN-CT-2003-504712, MIUR-PRIN 2004, MIUR-FIRB 2001, CrdC-AMRA, INFM-PCI.

<sup>†</sup> Electronic address: coniglio@na.infn.it

<sup>‡</sup> Electronic address: mario.nicodemi@na.infn.it; URL: <http://smcs.na.infn.it>

- [1] S. Chandrasekhar, Hydrodynamic and Hydromagnetic stability. (Clarendon, Oxford, 1961).
- [2] C.D. Andereck, S.S. Liu and H.L. Swinney, J. Fluid Mech. **164**, 155-183 (1986).
- [3] S.L. Conway, T. Shinbrot, B.J. Glasser, Nature **431**, 433-437 (2004).
- [4] T. Shinbrot and F.J. Muzzio, Nature **410**, 251-258 (2001).
- [5] H.M. Jaeger, S.R. Nagel and R.P. Behringer, Rev. Mod. Phys. **68**, 1259 (1996).
- [6] D.J. Goldfarb, B.J. Glasser and T. Shinbrot, T., Nature **415**, 302-305 (2002).
- [7] For general references on segregation see: J.M. Ottino and D.V. Khakhar, Ann. Rev. Fluid Mech. **32**, 55 (2000); A. Kudrolli, Rep. Prog. Phys. **67**, 209 (2004); F. Canteleube and D. Bideau, Europhys. Lett. **30**, 133 (1995); H.A. Makse, S. Havlin, P.R. King, and H.E. Stanley, Nature **386**, 379 (1997).
- [8] T. Mullin, Phys. Rev. Lett. **84**, 4741 (2000).
- [9] P.M. Reis and T. Mullin, Phys. Rev. Lett. **89**, 244301 (2002).
- [10] P. Sánchez, M.R. Swift, and P.J. King, Phys. Rev. Lett. **93**, 184302 (2004).
- [11] P.M. Reis, G. Ehrhardt, A. Stephenson and T. Mullin, Europhys. Lett. **66**, 357 (2004).
- [12] H.J. Herrmann and S. Luding, Cont. Mech. Thermod. **10**, 189-231 (1998).
- [13] G. Ehrhardt, A. Stephenson and P.M. Reis, cond-mat/0403273.
- [14] L.E. Silbert, D. Ertas, G.S. Grest, T.C. Halsey, D. Levine and S.J. Plimpton, Phys Rev E **64**, 051302 (2001).
- [15] P.A. Cundall and O.D.L. Strack, Geotechnique **29**, 47-65 (1979).
- [16] In colloidal systems the formation of stripes parallel to the driving direction has been numerically observed in the presence of a constant shear between the two components, as in J. Dzubiella, G.P. Hoffmann and H. Lowen, Phys. Rev. E **65**, 021402 (2002).
- [17] M.A. Scherer, F. Melo, M. Marder, Phys. of Fluids **11**, 58-67 (1999).
- [18] C.M. Pooley, and J.M. Yeomans, Phys. Rev. Lett. **93**, 118001 (2004).
- [19] R. Wunenburger, D. Beysens, C. Chabot, Y. Garrabos, P. Evesque and S. Fauve, Proceedings of the STAIF 2000, Albuquerque, New Mexico, USA.
- [20] A. Stegner, J.E. Wesfreid, Phys. Rev. E **60**, R3487 (1999).
- [21] If both species are monodisperse we do observe a phase diagram which is similar to that of Fig 3. However when segregated in stripes both species can be found in a liquid or in a crystal-like configuration.
- [22] “Unifying concepts in granular media and glasses”, (Elsevier Amsterdam, 2004), Eds A. Coniglio, A. Fierro, H.J. Herrmann, M. Nicodemi, P. Richard, M. Nicodemi, R. Delannay, P. Ribière, D. Bideau, Nature Materials **4**, 121 (2005).

\* Electronic address: picciam@na.infn.it